

**City of Falcon Heights
2077 W Larpenteur Avenue
Falcon Heights MN 55113**

**CITY COUNCIL
WORKSHOP AGENDA**

**February 4, 2009
6:30 p.m.**

- 1. Review of Revised Larpenteur Avenue Streetscape Proposal.**
- 2. Boulevard Tree Maintenance Program (See attached items).**

TO: Mayor Lindstrom, Council members Harris, Kuettel, Long, and Mercer-Taylor

FROM: Justin Miller, City Administrator

Re: Revised Larpenteur Streetscape Design Proposals

Explanation:

Over the past year, the City has been working with SRF Consulting to develop streetscape designs for Larpenteur Avenue. Initial work included soliciting input from the public and an advisory committee, and in late 2008 the consultants presented their findings to the city council.

Once the city council reviewed the entire plan, staff and the consultants were directed to return with a focus on addressing key intersections along the corridor and median plantings. Attached to this report are estimates for these two items.

Crosswalks (Figure A):

SRF has broken down cost estimates for each intersection along the Larpenteur corridor. If each intersection had the existing concrete removed and replaced with colored concrete, it would cost an estimated \$336,055. SRF investigated a less costly concrete staining product, but after discussions with Ramsey County staff it was determined that the application method would not be allowed and would not be as cost-effective.

The council could choose to do the work at each intersection or choose individual intersections at which to incorporate the colored concrete.

Median Plantings (Figure B):

Estimates for planting trees and landscaping at each major median were also prepared by SRF. If each major intersection were selected, this portion of the project would cost \$417,208.20. This would include concrete removal, installation of trees and sod, and other landscaping features. Again, the city council could choose to target these improvements at specific medians.

Staff and SRF personnel will be on hand to answer any questions as well as present more detailed funding source information at the workshop on Wednesday.

Figure: A CROSSWALKS Cost Breakdown	UNIT	UNIT PRICE	EST. QUANTITY	EST. AMOUNT
Fulham (1)				
Sawing Concrete Pavement	LF	\$2.00	80	\$160.00
Remove Concrete Pavement	SF	\$3.00	1200	\$3,600.00
Colored Concrete	SF	\$12.00	1200	\$14,400.00
Subtotal				\$18,160.00
Coffman Area (2)				
Sawing Concrete Pavement	LF	\$2.00	160	\$320.00
Remove Concrete Pavement	SF	\$3.00	1600	\$4,800.00
Colored Concrete	SF	\$12.00	1600	\$19,200.00
Subtotal				\$24,320.00
Cleveland (4)				
Sawing Concrete Pavement	LF	\$2.00	300	\$600.00
Remove Concrete Pavement	SF	\$3.00	4500	\$13,500.00
Colored Concrete (see note A)	SF	\$12.00	4500	\$54,000.00
Subtotal				\$68,100.00
Gortner (4)				
Sawing Concrete Pavement	LF	\$2.00	260	\$520.00
Remove Concrete Pavement	SF	\$3.00	2600	\$7,800.00
Colored Concrete (see note A)	SF	\$12.00	2600	\$31,200.00
Subtotal				\$39,520.00
Fairview (3)				
Sawing Concrete Pavement	LF	\$2.00	225	\$450.00
Remove Concrete Pavement	SF	\$3.00	3375	\$10,125.00
Colored Concrete (see note A)	SF	\$12.00	3375	\$40,500.00
Subtotal				\$51,075.00
Snelling (4)				
Sawing Concrete Pavement	LF	\$2.00	400	\$800.00
Remove Concrete Pavement	SF	\$3.00	6000	\$18,000.00
Colored Concrete (see note A)	SF	\$12.00	6000	\$72,000.00
Subtotal				\$90,800.00
Arona (4)				
Sawing Concrete Pavement	LF	\$2.00	220	\$440.00
Remove Concrete Pavement	SF	\$3.00	2200	\$6,600.00
Colored Concrete (see note A)	SF	\$12.00	2200	\$26,400.00
Subtotal				\$33,440.00
Hamline (1)				
Sawing Concrete Pavement	LF	\$2.00	70	\$140.00
Remove Concrete Pavement	SF	\$3.00	700	\$2,100.00
Colored Concrete (see note A)	SF	\$12.00	700	\$8,400.00
Subtotal Removals				\$10,640.00

Total Est. Amount **\$336,055.00**
Total Sq.Ft. **22175**

Note A: Colored concrete unit price assumes color added to full depth and pavement thickness less than 7". Pavement thickness greater than 7" will be slightly more.

Figure B: MEDIAN Cost Breakdown	UNIT	UNIT PRICE	EST. QUANTITY	EST. AMOUNT
Fulham to Coffman				
Remove Concrete Pavement	SF	\$3.00	8120	\$24,360.00
Colored Concrete	SF	\$6.00	1330	\$7,980.00
Streetscape Tree	TREE	\$325.00	16	\$5,200.00
Perennial planting (Native grasses and/or wildflowers)	PLT	\$12.00	1500	\$18,000.00
Sod Type Salt Resistant	SY	\$3.00	104	\$312.00
Landscape Edger	LF	\$4.45	40	\$178.00
Select Topsoil Borrow	CY	\$38.00	115	\$4,370.00
Shredded Bark Mulch	CY	\$25.00	15	\$375.00
Subtotal				\$36,415.00

Coffman to Cleveland				
Remove Concrete Pavement	SF	\$3.00	5150	\$15,450.00
Colored Concrete	SF	\$6.00	4600	\$27,600.00
Streetscape Tree	TREE	\$325.00	1	\$325.00
Perennial planting (Native grasses and/or wildflowers)	PLT	\$12.00	0	\$0.00
Sod Type Salt Resistant	SY	\$3.00	55	\$165.00
Landscape Edger	LF	\$4.45	20	\$89.00
Select Topsoil Borrow	CY	\$38.00	6	\$228.00
Shredded Bark Mulch	CY	\$25.00	4.5	\$112.50
Subtotal				\$43,969.50

Cleveland to Gortner				
Remove Concrete Pavement	SF	\$3.00	6830	\$20,490.00
Colored Concrete	SF	\$6.00	2630	\$15,780.00
Streetscape Tree	TREE	\$325.00	2	\$650.00
Perennial planting (Native grasses and/or wildflowers)	PLT	\$12.00	1875	\$22,500.00
Sod Type Salt Resistant	SY	\$3.00	88	\$264.00
Landscape Edger	LF	\$4.45	80	\$356.00
Select Topsoil Borrow	CY	\$38.00	17	\$646.00
Shredded Bark Mulch	CY	\$25.00	7	\$175.00
Subtotal				\$60,861.00

Gortner to Fairview				
Remove Concrete Pavement	SF	\$3.00	5220	\$15,660.00
Colored Concrete	SF	\$6.00	2320	\$13,920.00
Streetscape Tree	TREE	\$325.00	8	\$2,600.00
Perennial planting (Native grasses and/or wildflowers)	PLT	\$12.00	31	\$372.00
Sod Type Salt Resistant	SY	\$3.00	116	\$348.00
Landscape Edger	LF	\$4.45	40	\$178.00
Select Topsoil Borrow	CY	\$38.00	14	\$532.00
Shredded Bark Mulch	CY	\$25.00	4	\$100.00
Subtotal				\$33,710.00

Fairview to State Fair Entry				
Remove Concrete Pavement	SF	\$3.00	7870	\$23,610.00
Colored Concrete	SF	\$6.00	1670	\$10,020.00
Streetscape Tree	TREE	\$325.00	20	\$6,500.00
Perennial planting (Native grasses and/or wildflowers)	PLT	\$12.00	1250	\$15,000.00
Sod Type Salt Resistant	SY	\$3.00	488	\$1,464.00
Landscape Edger	LF	\$4.45	30	\$133.50
Select Topsoil Borrow	CY	\$38.00	124	\$4,712.00
Shredded Bark Mulch	CY	\$25.00	23	\$575.00
Subtotal				\$62,014.50

State Fair Entry to Arona				
Remove Concrete Pavement	SF	\$3.00	4750	\$14,250.00
Colored Concrete	SF	\$6.00	4750	\$28,500.00
Streetscape Tree	TREE	\$325.00	0	\$0.00
Perennial planting (Native grasses and/or wildflowers)	PLT	\$12.00	0	\$0.00
Sod Type Salt Resistant	SY	\$3.00	0	\$0.00
Landscape Edger	LF	\$4.45	0	\$0.00
Select Topsoil Borrow	CY	\$38.00	0	\$0.00
Shredded Bark Mulch	CY	\$25.00	0	\$0.00
Subtotal				\$42,750.00

Snelling Medians (north and south of Larpenteur)				
Remove Concrete Pavement	SF	\$3.00	10320	\$30,960.00
Colored Concrete	SF	\$6.00	2800	\$16,800.00
Streetscape Tree	TREE	\$325.00	10	\$3,250.00
Perennial planting (Native grasses and/or wildflowers)	PLT	\$12.00	0	\$0.00
Sod Type Salt Resistant	SY	\$3.00	782	\$2,346.00
Landscape Edger	LF	\$4.45	55	\$244.75
Select Topsoil Borrow	CY	\$38.00	86	\$3,268.00
Shredded Bark Mulch	CY	\$25.00	14.25	\$356.25
Subtotal				\$57,225.00

Arona to Hamline				
Remove Concrete Pavement	SF	\$3.00	9150	\$27,450.00
Colored Concrete	SF	\$6.00	8150	\$48,900.00
Streetscape Tree	TREE	\$325.00	3	\$975.00
Perennial planting (Native grasses and/or wildflowers)	PLT	\$12.00	150	\$1,800.00
Sod Type Salt Resistant	SY	\$3.00	44	\$132.00
Landscape Edger	LF	\$4.45	16	\$71.20
Select Topsoil Borrow	CY	\$38.00	20	\$760.00
Shredded Bark Mulch	CY	\$25.00	7	\$175.00
Subtotal				\$80,263.20

**Total
Est.
Amount \$417,208.20**

TO: Mayor Lindstrom, Council members Harris, Kuettel, Long, and Mercer-Taylor

**FROM: Justin Miller, City Administrator
Tim Pittman, Parks and Public Works Director
Andy Hovlund, City Forester**

Re: Boulevard Tree Maintenance Program

One of the 2009 City Council goals is to explore a boulevard tree maintenance program. Current practice calls for the city to install, at city expense, a boulevard tree on each residential property if one is requested. Likewise, if the tree is diseased or damaged during a storm, the city pays to remove and replace it. Any maintenance outside of routine pruning is the responsibility of the homeowner.

Periodically the city receives requests from homeowners who wish to chemically treat boulevard trees to prevent diseases such as Dutch Elm disease or Oak Wilt. The city has usually agreed to these requests, pending a signed release from the homeowner which lets the homeowner know that the city reserves the right to remove the tree if the city forester deems it to meet the disease criteria set forth by state statute.

Our city forester recently contacted several tree care companies about providing a city-wide quote for treatment of boulevard trees for diseases that can be chemically treated. He received two responses, one from S&S Tree Specialist, and the other from Rainbow Tree Care. Their quotes for elm treatments are below:

S&S Tree Specialist	\$8.95/diameter inch
---------------------	----------------------

Rainbow Tree Care	
Arbotech product	\$13/diameter inch
Alamo product	\$10.50/diameter inch

Both of the companies above use different products with varying degrees of success. Instead of staff attempting to select the best product, we are recommending that residents make their own decisions for what is best for their needs. The selected company could then bill either the city or resident, and the city would simply facilitate the process by securing the lower bid price from the above companies. We would still require a signed release acknowledging that the city reserves the right to remove the tree if the city forester determines it to be a hazard, and the city will not reimburse the

homeowner for any treatment costs incurred. Furthermore, staff is recommending that the city not invest any city funds into this project at this time due to budgetary constraints.

Our boulevard trees are a valuable asset to our community, and the city goes through great expense to provide this amenity. Further, they add value to our housing stock, so staff believes it is appropriate for our homeowners to share in the costs of maintaining this asset.

Staff and the city forester will be present at Wednesday's workshop to better answer any questions.

Protection from

Dutch Elm Disease

with MACRO-INFUSION™



- Macro-Infusion™ with Arbotect® 20-S for Dutch Elm Disease prevention is 99.5% effective for 3 seasons.
- Recommended by most major university's such as the University of Minnesota, University of Michigan, Kansas State University, University of Wisconsin and other major universities.
- Backed by 20 years of field use and research
- Used by leading arborist companies such as Davey Tree, Bartlett Tree Experts and The Care of Trees.



Arbotect® 20-S

is a systemic fungicide used in the treatment of Dutch Elm Disease and Sycamore Anthracnose by trained arborists and others trained in Macro-Infusion™ techniques.



Protection from
Dutch Elm Disease
with MACRO-INFUSION™

Table of Contents

Technical Description	pg.3
Symptoms and Diagnosis	pg.5
Managing The Disease	pg.6
Root Grafts	pg.8
Protecting Elms with Arbotect®-20S	
General Information	pg.9
Performing Macro-Infusion™	pg.10
Dosing Considerations	pg.13
Macro-Infusion vs. Micro Injection	pg.14
Saving Diseased Trees - Tracing	pg.15




Dutch Elm Disease

Technical Description


What is Dutch Elm Disease?

Dutch Elm Disease is a fungus called *Ophiostoma Novo-ulmi*. This fungus grows only in elms. Most species of elms are susceptible including, American, Slippery (red), English, European, and Winged. Less susceptible species include Siberian, Chinese, and Cedar elms. The disease is spread from infected trees to healthy trees most commonly on the elm bark beetle *Scolytus multistriatus*. It can also pass from one tree to another through root grafts, a situation where a tree's roots fuse underground with another tree of the same species.



The fungus *Ophiostoma Novo-ulmi* causes Dutch Elm disease. It is in the same fungus family as bread mold

Photo courtesy of University of Minnesota



Tracing can save elms if done in the initial stages of disease.

Fungal Growth Inside The Elm

The fungus that causes Dutch Elm Disease is not a blight or a wood decaying organism. Instead it lives inside the xylem of a susceptible elm. Understanding the fungus growth pattern is important to knowing if a infected elm can be saved and understanding the process of tracing, a method where an arborist creates a physical separation of the fungus from the elm.

Beetle infections generally start in the 2 - 4 year old twigs. This is where the beetle feeds and mates. The fungus rubs off the beetle and begins to grow in the tree. The fungus grows in a very predictable pattern. Usually it stays narrow in width and grows down the branch. It will enter the stem of the tree and grow in this narrow line to the roots. Once the fungus enters the root flares, the fungus will grow sideways and back up the tree in many places. Tracing can save elms if the initial infection is caught before entering the roots.

A characteristic stain on the xylem of an elm infected with Dutch Elm Disease is caused by the tree producing gum like substances called tyloses in an attempt to stop the spread of the disease. These tyloses actually are what cause the tree to wilt and die as they block the xylem and prevent water transport to the top of the tree. So in effect the fungus stimulates the elm to kill itself.




Macro-Infusion of Arbotect® protects healthy Elms for 3 growing seasons and is 99.5% effective.

Dutch Elm Disease

Technical Description

The Beetle That Spreads Dutch Elm Disease

An elm bark beetle's life revolves around elm trees. It breeds in dead and dying elms, and depends exclusively on elm tissue for food, this creates a devastating cycle for elms, when the Dutch elm disease fungus *Ophiostoma nova ulmi* becomes part of the cycle. Thousands of eggs can be laid by the adult females in one piece of wood the size of a fireplace log. If the tree died from Dutch Elm Disease every beetle that hatches and emerges as an adult from that wood may be carrying the fungus. They will be looking for a healthy elm on which to feed and will inadvertently inoculate the tree with Dutch Elm Disease via the spores carried on their bodies. The majority of new infections, and the bulk of the losses to Dutch Elm Disease, are due to this method of transmission. That is why prompt detection, removal, and destruction of dying elms is so critically important.



This wound on this small 4 year old twig was caused by a beetle that introduced Dutch Elm disease to this tree.



Photo courtesy of University of Minnesota

The Beetle *Scolytus multistriatus* spreads the disease from infected to healthy Elms.



healthy tree

infected tree



Root grafts can also spread the disease from sick to neighboring healthy elms

Photo courtesy of University of Minnesota

Root Graft Spread of Dutch Elm Disease

The other method of disease transmission between elms is through grafted roots. When elms are growing near each other their roots come in contact in the soil and graft together. In the absence of a vascular wilt disease, such as Dutch Elm Disease, this is an advantage for the trees. The Dutch elm disease fungus however can pass from diseased to healthy trees through these grafted roots and continue to spread indefinitely through a stand of elms.


Dutch Elm Disease

Symptoms & Diagnosis

Symptoms

The first evidence of Dutch Elm Disease is wilting or "flagging" in one or more of the branches, usually starting in the outer portion of the crown. Leaves on the infected branches turn dull green to yellow and curl, finally becoming dry, brittle, and brown. The symptoms progress down the limb and eventually throughout the entire tree. Trees infected through root grafts can die very rapidly, while trees infected via the feeding of bark beetles can take 1 - 3 years to die. Another symptom of the disease is the discoloration of the water conducting vessels. This is easily seen by peeling the bark off infected wood, revealing the brownish staining.

Foliar symptoms of Dutch Elm Disease



Diagnosis

Understanding the Dutch elm fungus and how it grows makes diagnosing this disease easier. A few key distinctions will help in accurately making sure that you are in fact dealing with this dreaded disease.

1. Leaves on infected trees are almost always curled or wilted looking. They often drop off. Leaves that are flat and shiny are common in the late summer and are usually caused by branch senescence.
2. The disease symptoms progress as the fungus grows in the tree. Thus, there is a pattern of leaf death outside the tree that reflects the fungus growth inside the tree. Remember that the initial infection grows downward in a narrow band until it has reached the roots. It then spreads sideways and grows back up the rest of the tree. See the Technical Description of Dutch elm disease for more information.
3. Check under the bark close to where there are external leaf symptoms. Dutch Elm Disease always causes the water conducting vessels to turn a dark brown. Finding this discoloration along with wilting leaves is a very good indicator that Dutch elm disease is present. Use a chisel and a hammer to open a hole in the bark to check for the discoloration
4. Trees in the spring can die rapidly – causing the appearance that the disease is moving very quickly. What is happening in reality is the fungus was in the tree from last summer, the tree grew new tissue over the top of the infected wood, and then the fungus colonized the newly formed wood causing rapid dieback. Infections that happen in the current year can be seen reflected in leaf dieback as the fungus grows.
5. If you are unsure if your elm has Dutch Elm Disease– you can take samples and send them to a lab for confirmation. See sending samples to a lab for diagnosis.

Another symptom is discolored brown to black wood under the bark. Healthy elm wood is white.



The fungus grows downward to the roots. After reaching the root area, it spreads sideways and moves back up the tree.

Dutch Elm Disease

Managing the Disease

Dutch Elm Disease is difficult to control and without management it will wipe out a large population of elms in just a few years. However, with a properly implemented program, the devastating effects of the disease can be greatly reduced. An effective program includes four parts: prompt detection and removal of diseased elms, disruption of root graft transmission, saving diseased elms with tracing, and protection of valuable elms through proper Macro-infusion procedures.



Before - A beautiful property before Dutch elm disease



After - The same property after the devastation of Dutch elm disease.



Photo courtesy of City of Evanston

Scouting for Dutch Elm Disease (DED)

Scouting for Dutch Elm Disease and identifying diseased elms is the first step of any DED program. Scouting also involves checking people's yards and garages for elm wood. Scouts will usually move through an area every 2-4 weeks during the growing season to make sure dying trees are identified and properly dealt with.

Scouting is the first step of any Dutch elm control program

Removing Diseased Elms

Promptly removing and disposing of elms dying from Dutch Elm Disease is the key to effectively managing Dutch Elm Disease on a community wide basis. It involves identifying diseased elms that cannot be saved by tracing and immediate removal of these trees. This will reduce the number of disease carrying beetles. A single dead elm can produce tens of thousands of contaminated beetles. Without such a program, a substantial majority of a community's elm population will die within a few years. Removed elms need to either have their bark removed, or be chipped, burned, or buried. Storage of diseased elm wood must not be anywhere near valuable elm populations.




Dutch Elm Disease

Managing the Disease

Root Graft Disruption

Severing grafted root systems between diseased and healthy elms can save many trees. This is most reliably accomplished mechanically with either a trencher or vibratory plow. In areas where there are utilities an air spade can be used to open a trench so that root grafts can be cut without damaging the utility lines. Remove the diseased elm only after common roots have been disrupted. Accordingly, prompt disease detection as well as installing root graft barrier trenches is very important to the overall success of an integrated control program. See section on root graft disruption.



Saving Infected Elms with Tracing

Tracing is a method of saving recently infected elms. It is far more cost effective than removing and replacing an elm. By utilizing this procedure, a city can save many thousands of dollars. Since the fungus that causes Dutch Elm Disease grows very predictably in a narrow band downward and only in the current years water conducting vessels – this methodology is easy to learn and incorporate. See section on saving diseased elms with tracing.



Root grafts must be physically severed to prevent spread of Dutch elm from tree to tree through the roots.



Before - This tree was traced and saved



After - This is the same tree 6 years later. Alive and doing well.



Macro-infusion of Arbotect protects elms for 3 growing seasons from Dutch Elm Disease and is 99.5% effective.

Protecting Valuable Individual Trees

There have been 100's of proposed "cures" or protection products for Dutch Elm Disease over the years, but only one treatment has been proven to work through replicated University trials - macro-Infusion with Arbotect 20-S. Arbotect gives greater than 99% success over 3 years when applied properly. Arbotect does not work if the tree already is infected with Dutch elm disease or the disease fungus enters the tree through root grafts.

Dutch Elm Disease

Root Grafts

When elms are growing near each other their roots come in contact with one another in the soil and graft together. In the absence of Dutch Elm Disease this is an advantage for the trees. The Dutch Elm Disease fungus, however, can pass from diseased to healthy trees through these grafted roots and continue to spread indefinitely through a stand of elms.

Physical Breaking of Roots

The only way to stop the spread of Dutch elm disease through root grafts is by physically breaking the root connections between the infected tree and the healthy tree.

This is most often accomplished with trenchers or vibratory plows. In situations where there are buried utilities, an Air Spade™ or AirKnife™ can be used to remove the soil and expose the root grafts without damaging the utility lines. Trenches should be at least 3' deep in clay soils and 5' in sandy soils, although it is always best to go as deep as your equipment allows. .



Install Trenches Midway Between Infected and Healthy Tree

Trenches are typically installed midway between the infected tree and the healthy tree, although this can be adjusted in either direction depending on how far the infection has spread.

It is important to know where the fungus is located in the diseased tree before the trench is installed. If the disease stain is already at ground level in the infected tree it is impossible to determine how far it has traveled through the root system toward the healthy tree, so a second trench may be necessary. This trench would isolate the healthy appearing tree from any others that might be root grafted to it.

Sever the Root Grafts Before Removing the Infected Tree

It is important to sever the root grafts before removing the diseased tree. Because each tree is transpiring, removal of the diseased tree must allow the healthy tree to pull all of the moisture and fungal inoculum out of the other tree's root system quickly.



An airspade can be used to remove soil and expose root grafts in situations where utilities are present.



Elms that grow close to each other (Canopies touching) are often root grafted.



Root grafts must be physically broken to prevent Dutch Elm Disease from growing into a healthy elm from a diseased elm.

Dutch Elm Disease

Macro-Infusion™ General Information

Arbotect Lasts 3 growing seasons

Arbotect is a systemic fungicide that is highly resistant to degradation and moves into new sapwood in sufficient quantity to protect trees for 3 growing seasons.


The Purpose Is Complete Distribution

The purpose of preventive macro-infusion with Arbotect is to provide even and complete distribution of the fungicide throughout the 2-4 year old twigs where the beetles feed.



Rainbow Treecare has had a 99.5% success rate on a group of about 6000 trees under protection over a 3 year period.

Macro-infusion of Arbotect protects Elms for 3 growing seasons from Dutch Elm Disease.



The purpose of macro infusion is to evenly and completely cover the entire internal crown of the tree with enough Arbotect to keep the disease out.

How Long It Takes

The average uptake for 20-60 gallons of fungicide solution is 30-60 minutes if the protocol is followed closely. Trees should be treated after the leaves have fully enlarged or after the seeds have dropped, and the treatments can be administered until there is fall color present in the canopy.

Arbotect Limitations

Diseased elms should NOT be treated with Arbotect. Rainbow's experience is that Arbotect will mask the symptoms of the disease for 1-3 seasons before the tree dies. If your elm is already infected with Dutch elm disease, tracing may be a method that can be used to save the tree. Arbotect does not prevent infections that grow through root grafts, so in situations where this is a concern, install a trench and wait until the following season to make sure the tree is healthy before protecting with Arbotect. As a preventive treatment this process gives highly predictable results.



Do not infuse symptomatic elms or elms that will become infected through root grafts. The process will not be effective.



Macro-Infusion can be done after the leaves are fully formed on the tree and until the tree begins to show fall coloration.



Treatment must be done into the root flares to get even coverage. Do not infuse into the trunk – this will result in poor distribution.

Dutch Elm Disease

Performing Macro-Infusion™

The Purpose Of Macro-Infusion

The purpose of Macro-Infusion is the complete and even coverage of the 2 - 4 year old branches so that if a beetle comes to feed, the disease cannot enter the tree. Infusing into the root flares is critical to the success of even and complete distribution. The tissue in the root flares allows for lateral movement of the chemical as it is infused. Trunk tissue on the other hand is hard and rigid. It is highly compartmentalized. Infusion into trunk tissue will provide very limited lateral movement of the Arbotect solution and will result in small portions of the crown receiving a large dose of Arbotect with other areas receiving nothing.

The Process

Step 1 - Inspect the tree

- Make sure the elm you are treating is not diseased. Fully scan the crown to make sure there are no symptoms. If the tree is diseased - DO NOT TREAT. Use the tracing process to isolate any disease that might be present.
- Measure the tree and determine your dosage. Elms require 12 oz of Arbotect per 5" DBH. (Diameter breast height). Each ounce of Arbotect is diluted in 2 gallon of water. You can find a copy of the DOSAGE TABLE by clicking on the link.
- Large trees could require increased amounts of chemical to achieve complete distribution. Small elms, or trees with portions of the crown missing may require a reduced dose. See Dosing Considerations for more information.
- Determine how much root flare excavation is needed.
- Look for root rot or significant decay in the root collar area - if present do not treat.

Elms infected with Dutch elm disease should not be treated

Significant canopy die back or stress may compromise uptake time and distribution and be a sign of a serious problem slowly killing the tree

Step 2 - Excavate the root flares (if necessary)

- Use a shovel and trowel to remove sod and soil without damage to the tree
- Thoroughly brush soil from root flares with a hand broom
- Infusion sites should be 8-10 inches below the top of the root flare

Soil left on the root flare can dull the bit and plug the xylem.

If sod is carefully removed it can facilitate fast cleanup.



Remove all soil from root flares with a brush. Drilling into a dirty flare will dull your bit and push dirt into the hole – slowing down the uptake.



Equipment need to perform Macro-Infusion on elms.



Every aspect of macro infusion is designed to get even and complete crown coverage of the Arbotect solution.



Infusing into the root flares is critical to achieve even and complete distribution of the Arbotect solution throughout the crown of the tree.

Dutch Elm Disease

Performing Macro-Infusion™

Step 3 - Drilling the holes

- Use a sharp, high helix drill bit (change every 5 trees)
- Drill perpendicular to the surface of the flare
- Drill one inch past the bark
- Drill at slow speeds and do not excessively spin the bit in the hole
- Use 1 to 1-1/2 infusion sites (in the root flare) per diameter inch (measured at breast height)
- Place at least one infusion site on each root flare

Do not place infusion sites into or below dead tissue

Do not drill into deep valleys or sunken areas

Begin filling the reservoir with water while drilling the holes



Drill a series of small one inch deep holes around the tree at the root flares.

Step 4 - Inserting the tees

- The current year xylem are the only vessels that will take up the solution. Make sure your tee is properly positioned to deliver the Arbotect there.
- Check each tee to be sure it is not plugged and replace any that are badly damaged
- Firmly insert tees by hand and very lightly tap each tee to set it
- Attach tubing from solution reservoir to feed into the harness in 2 locations. These 2 sites should be on opposite sides of tree

Plugged tees will prevent that portion of the xylem from receiving chemical

The only vascular tissue that conducts water is the current year's xylem



Use a sharp HIGH HELIX drill bit for fastest uptake. Replace drill bit every 5 trees.



Insert infusion tees and hook up tubing harness to pump.



Make sure the Macro - Infusion tee is delivering solution to the current year's xylem. This is the only water conducting tissue that will take up solution.

Step 5 - Starting the Infusion

- Pull out 2 tees on opposite sides of the tree
- Prime the pump (if it is not self-priming)
- Turn on the pump and bleed the air out of the line
- With all air out of harness, re-insert the 2 tees and check for leaks
- Adjust the pressure to 15-20 psi
- Lightly tap any leaking tees

Increasing the pressure will not make the infusion go faster

If a tee persists in leaking, drill a new hole.

Use a very small hammer to tap leaking tees - this helps prevent driving them in too far

Dutch Elm Disease

Performing Macro-Infusion™

Step 6 - Mixing the Chemical

- Do not mix the chemical into the reservoir until the system is running and free of leaks
- Arbotect will form a white precipitate in water that is hard or has high pH.
- Mix a small amount of Arbotect and water in a clear measuring cup
 - If the solution stays clear add the appropriate dose to the reservoir
 - If a white precipitate forms and falls to the bottom of the cup, the water needs to be treated
 - Add muriatic acid to the reservoir at 1 oz. per 6 gallons of water
 - Add the Arbotect and watch for white precipitation
 - If no precipitate forms, continue with the infusion
 - If the precipitate forms again, continue to add muriatic acid at the rate listed above until the Arbotect stays in solution
- Passing water through a deionizing tank also prevents Arbotect precipitation problems



Be careful about water source. Treat water with muriatic acid if pH is high. Avoid hard water or run through a deionizer. Arbotect will precipitate out of solution in hard and high pH water

Muriatic acid is available at hardware or building supply stores.

For infusions that require a larger volume of solution than the reservoir holds, use 5 gallon buckets to measure the remaining dose and add to the reservoir.

White air bubbles may be seen floating in the reservoir if air gets into the lines. Do not confuse this with when Arbotect precipitate falls out of solution to the bottom of the reservoir.

Step 7 - During the Infusion

- Monitor tees for leaks
- Maintain pressure at 15-20 psi
- Pack other equipment such as drill, unused chemical, etc.
- Prepare other trees on site for treatment



Macro-Infusion uptake should take less than 1 hour – 90% of the time. Call Rainbow for technical support if uptake is consistently longer than this.

Step 8 - Cleanup

- Turn off pump when air is drawn into the harness
- Remove tees from the tree
- Replace soil and sod around base of tree

Do not plug the infusion holes with wax, wooden plugs, or any other substance

Rainbow Treecare uses 30 gallon trash cans for the solution reservoir and packs all equipment in these when the infusion is over



Macro infusion is 99.5% successful in protecting elms for 3 years when performed properly.

Dutch Elm Disease

Dosing Considerations

Dosing Considerations When Performing Macro-Infusions

The purpose of macro-infusion is to provide enough chemical so that it is evenly distributed throughout the entire crown of the tree. If the dose is too high there may be a phytotoxic reaction, and if the dose is too low infections may occur.

Dosage is determined by measuring the diameter of the trunk at breast height (DBH). Trunk diameter alone, however, does not always reflect the proper dose because it may not reflect canopy development. Very few trees are open-grown in the urban setting. Their canopy development has been influenced by the proximity of buildings, other trees, power line clearance, etc. Our experience in treating trees over the past 20 years has taught us that proper dosing is an art as well as a science, and it is a key to predictable results.

How to Dose Trees

1. Use the label rate for trees between 22" – 28" DBH with average size crown and no major limbs missing. For smaller trees, reduce the dose by 3% for every inch below 20" DBH.
2. For trees with significant canopy losses, it is necessary to reduce the dose accordingly. For example, if 15% of the canopy was lost due to storm damage, reduce the dose by 15%.

<p>Example #1 A 25" DBH elm with approximately 30% missing crown from pruning and storm damage:</p> <p>Step 1 - Label rate = 12 oz per 5" DBH = $25"/5" \times 12\text{oz} = 60\text{oz}$</p> <p>Step 2 - Estimate of missing crown = Subtract 30%</p> <p>Total dose = 60oz – 30% (18oz) = 42oz Arbotect</p>	<p>Example #2 A 17" DBH elm with a full crown</p> <p>Step 1 - Label rate = 12oz per 5" DBH = $17"/5" \times 12\text{oz} = 41\text{oz}$ Subtract 3% per inch under 20" = sub ~ 9%</p> <p>Step 2 - Estimate of missing crown = Minus ~ 0%</p> <p>Total dose = 41 oz - 9% (4 oz) = 37oz</p>
---	---

3. Large trees (above 28" DBH) require a higher dose to achieve the best results. Increasing the dose above the label rate, however, is not legal and we do not recommend you break the law. Rainbow Treecare is conducting research to determine optimal doses for larger trees and will use these results to make appropriate label changes. The procedure we will test for large trees, consists of making reductions to the canopy, as described above, and increasing the dose by 3% for every inch of diameter above 28" DBH.

Symptoms of Overdose

Overdoses tend to occur on smaller trees, and the effects are temporary for the tree. When overdosed, leaves tend to turn a gray brown color and may curl or remain flat. Affected leaves may drop from the tree. Symptoms tend to develop in the lower canopy, and often on suckers of elm trees. Damaged trees may re-leaf later in the season or bounce back the following spring.

Slippery elms (*Ulmus rubra*)

Reduce the dose to one-half of the label rate. A full dose may give phytotoxicity.

Macro-Infusion™

Advantage

Complete Crown Coverage with Macro-Infusion

Macro-Infusion is a large volume application of fungicide/water solution. Research shows that fungicides need a high volumes of a carrier to get complete distribution throughout trees.

Fungicide Macro-Infusion vs Micro-Injection

	Macro-Infusion	Micro-Injection
Chemical Carrier	Water	Alcohol/Xylene
Years Between Treatment	3 years	1 year
Tree Injury	Low	High
Volume of Material	20 - 40 gallons	1-2 ounces
Distribution in Tree	Complete	Minimal
Application Time	40 - 90 minutes	30 to 40 minutes
Injectio Wounding Size	15/64"	11/64" *

*Some micro-injectors use smaller holes



Proven Fungicides

Macro-Infusion™ of Arbotect and Alamo is the treatment method recommended by universities for Oak Wilt and Dutch Elm Disease.

This includes:
Texas A&M
Texas Forest Service
University of Minnesota
Michigan State

Complete Distribution Requires Root Flare Application

The science of infusing materials into trees has evolved. The purpose of any infusion is even and complete distribution of the material in the tree. This is difficult to accomplish with trunk injection. Trunk tissue is hard and compartmentalized preventing lateral chemical movement. Root flare injection allows lateral movement of material and complete crown coverage.

The Difference between Insecticides and Fungicides

There is a substantial difference in the mode of action between an insecticide and a fungicide. Insecticides work in small quantities and their mode of action is to kill the intended target, thus micro injection of insecticides can be effective. Many fungicides are really fungistatic in nature, this means they do not kill the fungus – but merely prevent it from growing. Once the fungistat chemical is gone – the fungus will continue to grow again. Fungistatic materials need to be administered in much higher volumes, this is why Macro-Infusion protocols are designed to deliver large quantities of fungistatic solution to get significant amounts of product into the crown of the tree.

Dutch Elm Disease

Saving Diseased Trees- Tracing

Macro Infusion Does NOT work on Diseased Elms

Our company has never had success saving a symptomatic elm infected with Dutch elm disease using Macro-Infusion. What typically happens when a diseased tree is treated with Macro-Infusion is the symptoms of the disease become masked and the tree then dies in 2 -3 years. We have had tremendous success utilizing a process we developed called tracing.

The 4 Distinctions you Need To Know to Successfully Save Infected Elms

In recent years developments have accrued which make saving infected elms a possibility. It is now possible to save elms that are infected into the trunk using a process that physically isolates the Dutch elm disease fungus. To understand this process you need to become familiar with 4 key distinctions.

Tracing can save elms diseased into the trunk.



Distinction 1 -

How the Fungus Grows

The Dutch Elm Disease fungus starts in the 2-4 year old twigs. It grows down the branch in a narrow band until it reaches the root system. Once in the root flares, the fungus can move laterally and spread rapidly throughout the tree. For Tracing to work, the fungus must not have entered the root flares.



This wound on this small 4 year old twig was caused by a beetle that introduced Dutch Elm Disease to this tree.



The fungus grows down the tree in a narrow band until it reaches the roots.


Dutch Elm Disease

Saving Diseased Trees- Tracing


Distinction 2

The Fungus Growth Is Predictable

The Dutch Elm Disease fungus is highly compartmentalized by the tree and grows in a very limited fashion. It can only grow in the current year's water conducting tissue, and has very limited lateral movement because the cell walls in the trunk and branches are very rigid. Thus, the fungus can be easily isolated from the rest of the elm by making some narrow and shallow cuts.



The fungus only grows in the current year's xylem.



By making some narrow and shallow cuts, the fungus can be isolated.

Distinction 3

The Fungus Can Only Grow Straight Down From A Branch Into The Trunk

Branch vascular tissue is only connected at its base to the trunk vascular tissue. This knowledge adds more predictability for the arborist practitioner on where the fungus is growing and where to look to begin the tracing process.



The branch vascular tissue is only connected to trunk vascular tissue at its base.

Distinction 4

The Stain is Produced By The Tree – Not The Fungus

The stain found under the bark is actually produced by the tree in response to the fungus and serves as your guide for the tracing process. It takes time for the stain to form and thus is not exactly representative of how far the fungus has grown down the tree. As a general rule, the stain will be 5 feet or so behind the fungus, thus the Arborist practitioner needs to continue the tracing process an additional 10 feet past the stain to ensure the fungus has been isolated.

Tools Needed

The tools needed include a small chain saw, a hammer, and a large flat headed screw driver. As well as the equipment to get up in the tree safely.



The stain under the bark is actually produced by the tree in response to the fungus. The fungus itself is growing ahead of the stain.

The Process of Tracing (PDF)– A detailed description of how to save elms with tracing.

× Rainbow Treecare
Scientific
Advancements

Advancing the Science of Tree Care
[About Us](#) | [Careers](#) | [Contact Us](#) | [Suggestions & Feedback](#) | [Site Map](#) |
[Links & Resources](#) | [Home](#)

Tree Health Issues

Bacterial Leaf Scorch

Chlorosis

Dutch Elm Disease

- [Technical Description](#)
- [Symptoms & Diagnosis](#)
- [Managing the Disease](#)
- [Root Graft Disruption](#)
- [Arbotect vs. Alamo](#)
- [Protect with Arbotect](#)
- [Arbotect MSDS](#)
- [Arbotect Label](#)
- [Saving Infected Elms](#)
- [Marketing Literature](#)
- [Research](#)
- [Library & Downloads](#)

Injured Roots

Insects

Oak Wilt Disease

Pine Wilt Disease

Reduce Growth

Sycamore Anthracnose

Urban Tree Stress

Vegetation Management

Products & Solutions

Alamo Fungicide

Arbotect Fungicide

Bacastat

Cambistat Growth

Regulator

Verdur

Xytect

Macro-Infusion

Micro-Infusion

Soil Injection

Marketing Materials

Research

Document Library

The following is the position of our Sister Company, Rainbow Treecare, on the use of Alamo instead of Arbotect for the treatment of Dutch Elm Disease. Rainbow Treecare Scientific Advancements fully supports the position of RTC and other top tree care companies around the country that Arbotect is the only fungicide with a proven, 30-year record of protecting elm trees. As the national marketer for both chemistries we are intimately familiar with the performance of each product. Alamo is very effective in the management of Oak Wilt Disease but experience and research shows it does not have three full years of efficacy in the prevention of DED as Arbotect does. Please [contact us](#) if you would like more information on these two products.

Rainbow Treecare Position Summary:

What Our Experience Tells Us about Arbotect and Alamo in Protecting Trees from Dutch Elm Disease

- Rainbow Treecare treats approximately 8000 elms on a 3-year protection cycle. We use Arbotect in the protection of those trees. Our loss rates average about 1%. Every treated elm is surveyed at least once, every three years, by one of our certified Arborists. That's an extra service many companies don't offer.
- Rainbow protects more trees from Dutch Elm disease than any company in the United States.
- The vast majority of companies treating elms in the U.S., who switched from using Arbotect to Alamo (propiconazole) in the late 1990's (with promises of results similar to Arbotect -- and faster on-site treatment times and lower costs) returned to Arbotect after substantial elm losses during the second and third years of the protection period.
- Rainbow Treecare treated 500 elms with Alamo (propiconazole) in the late 1990s in the Twin Cities and found that the treatments worked, but only for one year. Following those initial years we experienced major losses. This was very costly to our clients and

[Product Catalog \(PDF\)](#)

[MSDS/Labels](#)

Search our website

Newsletter Sign Up

enter email address:

1-877-ARBORIST

to our company. Please see [Wachtel](#) (in the Milwaukee area) and [Swingle Lawn, Tree, & Landscape Care](#) (in Denver), which experienced similar results ([Wachtel experience & Swingle experience](#)). Their [results](#) and [conclusions](#) are posted on our web site.


- Annual treatments are a possibility with Alamo (propiconazole) – but the issue of annual wounding (at the root flares for the injection process) and early season infection (pre-treatment) are reasons for concern. Given the wounding issue, and the fact that Alamo (propiconazole) does not move into annual new wood (necessary to give the tree the protection it requires), makes it clear to us that Alamo (propiconazole) fungicide does not offer a viable long term protection option.
- We have observed some therapeutic value with Alamo (propiconazole) during the first growing season following treatment. However, we have never actually experienced saving an infected tree using Alamo (propiconazole) (arresting the disease). Eventually, each time, the fungus has overcome and claimed the infected tree.
- By using Arbotect on a three season application cycle, via the macro-infusion method, we have protected literally thousands of elm trees from Dutch Elm disease across the Twin Cities area. The same has been true for other tree care companies across the United States who have adopted this same strategy.
- [Virginia Polytechnic Institute and State University](#) did a major study on Alamo (propiconazole) and its persistence. They could not find evidence of the chemical, at high dosage rates, in studies of treated trees eight months following injection. It was determined that Alamo (propiconazole) DID NOT move into the tree's new growth wood.
- Two other research papers have also been published by the [University of Minnesota](#) and [North Dakota State University](#), both recommending the use of Arbotect as the best chemical protection strategy against Dutch Elm disease.
- Two other published studies, by Mark Stennes (**Dutch Elm Chemotherapy with Arbotect 20-S and Alamo**) and R. Jay Stipes (**The Management of Dutch Elm Disease**) both support observations that Arbotect fungicide does in fact move into new wood (and offers multi-year protection), whereas no such observations have been documented for Alamo (propiconazole).
- We provide a three season treatment guarantee with Arbotect (treatment money back during the coverage

period if infection occurs), but only a one season guarantee with Alamo (propiconazole) (ask for a copy of these detailed guarantees from your Rainbow Arborist). Our first fungicidal recommendation, for uninfected trees, is always Arbotect.

- Some generic, Alamo-type chemistries have recently made their way into the U.S. market. We have nothing to do with these chemistries. We do not know that these products truly represent the chemistry found in propiconazole. These products are untried, untested, and uncontrolled. Therefore, we cannot support the use of these products in any way. Consumers approached to purchase these untested chemistries should be extremely wary.
- Rainbow Treecare offers mastery in the other (non-chemical) elements of DED management as well, including: neighborhood sanitation protocols, experienced tree climbers for whole tree inspections, isolation and tracing techniques, implications of root grafting, trenching and girdling expertise.
- Rainbow currently employs 30 ISA certified Arborists, with approximately 200 years of collective tree care experience that comes to bear when protecting valuable Twin Cities trees. Nearly all of Rainbow's arborists hold college degrees in forestry, urban forestry, or horticulture. Rainbow has the experience and the formal education to make recommendations that are based on a weight of hard facts, and not isolated conjecture.
- Rainbow Treecare's sister company, Scientific Advancements, is the sole distributor for both Arbotect and Alamo (propiconazole) fungicides (manufactured by Syngenta – formerly Novartis) to other tree care companies throughout the United States. We truly understand these chemistries and how they best fit within the context of a comprehensive battle against Dutch Elm disease. Our first recommendation, given the university studies and our years of our personal experience, is to select Arbotect as the # 1 choice in elm protection chemistries, as part of a comprehensive elm protection strategy. Alamo (propiconazole) is ideal for controlling Oak Wilt fungus, but not Dutch Elm disease.

Release Date: 06-05-06

© 2006. All Rights Reserved



Rainbow Treecare™
Scientific Advancements

2239 Edgewood Ave. S
St. Louis Park, MN 55426
1-877-ARBORIST
1-612-922-3810

Protocol:

Protecting Healthy Elms From Dutch Elm Disease For Three Seasons Using Arbotect 20-S

Protocol Information:

Predictable Results – This Process has been effectively used since 1980 on tens of thousands of elms. Rainbow Treecare has a 99.5% success rate in a three year injection cycle with a group of over 5500 elms. Untreated elms in the surrounding areas are lost at rates between 7.5% - 30% of the population. Results show a high degree of predictability.

Research – Completed at the University of Minnesota in 1979. Dr. David French and Mark Stennes 1987. *Distribution and Retention of Thiabendazole Hypophosphate and Carbendazim Phosphate Injected into Mature American Elms*. Phytopathology 77:707-712. G. N. Lanier 1987. *Fungicides for Dutch elm disease: Comparative evaluation of commercial products*. Journal of Arboriculture 13:189-195

Best Candidates for Treatment – High value American elms with no symptoms of Dutch elm disease and no root grafts to diseased trees. Good candidates are also free of root or crown rot diseases.

Treatment Method – Arbotect 20-S is macro-injected into the root flares of the tree with about 15-PSI pressure. Average uptake of material will be between ten minutes to one and a half hours after hook up. Occasionally a tree may take up to three hours or longer. Fastest uptake occurs on sunny mornings with a light breeze. The objective of this injection technique is to ensure the complete and even distribution of the chemical throughout the crown, enhancing the efficacy of the treatment. (For more information, please refer to the information sheet “How to Inject Trees Using Systemic Fungicides”). A three-year interval between treatments is important to minimize wounding in trees undergoing long term protection.

Dosing Considerations – Reduce label dosage for elms under 18” diameter or elms with reduced crown size. (See Rainbow information sheet “Dosing Considerations when Performing Macro Injection”)

Timing – Treat only after the leaves have reached full size and then throughout the summer. Do not treat if the tree has begun fall coloration. Delay treating trees with insect caused defoliation until the trees re-leaf.

Treating Elms Root Grafted to Other Diseased Elms – It is imperative to prune all disease out of the elm before treating. If the disease has progressed into the trunk, use the tracing method outlined in Rainbow information sheet “Tracing: A Physical Control for Dutch Elm Disease”. If the disease has grown into the root system, do not treat and remove the tree.

Side Effects – Injection wounds are one inch deep and 15/64” wide. Most elms compartmentalize well except those with root or crown rot diseases. In a small percentage of treated trees, some leaves in lower branches and suckers turn bronze and defoliate. Damage is usually temporary and the tree will re-leaf in a short time.

**City of Falcon Heights
2008 Year End Review-Forestry**

Trees Removed

Public

Diseased

Grove Park	18" elm behind playset
1597 Snelling Ave West	5" elm
1492 California	60" elm
2153 Hoyt- responsibility of U of M	30" elm

Gen. Decline

1547 Idaho Avenue	27" green ash
1423 Idaho	8" sugar maple
1724 W. Albert-severe decline and dieback exhibited	7" sugar maple
1525 Idaho-severe trunk decay in larger tree, hazardous	25" sugar maple
1477 California- tree in severe decline, leaning, suspect girdling roots	9" linden
1942 Summer-severe storm damage	21" red maple
1503 Hoyt-inspected for EAB, none detected	13" green ash
1872 Asbury-storm damaged/split during storm	14" green ash
1427 Iowa- poor specimen, low vigor	3" black ash

Declining sugar maples to remove and replace in spring
1379 Idaho, 1403 Iowa, 1406 Idaho, 1395 Idaho

Private

Diseased

2154 Folwell	22" elm	resolved
1830 Larpenteur	18, 13" elms-U of M	resolved
	5, 14" elms	resolved
1717 Larpenteur-Falcon Arms Apartmnts	9", diseased stump sprouts	resolved
1857 Fairview	7, 6, and 4" elms	
Falcon Heights Apartments	18" elm	resolved
2280 Folwell	7" elm	
1402 Idaho (along alleyway)	3" elm	resolved
1436 Roselawn	27" elm	resolved
1395 California (along alleyway)	3" elm	resolved

Insect

1596 Vincent-declining white oak 2-line chestnut borer (suspected)

Decline/Hazard

1771 Arona Street consulted with resident
storm damaged black walnut
resolved-applied to res propr taxes

Suspected DED Trees- Monitor in Spring and make determinations

1841 Pascal
2111 Hoyt
1564 Burton
1364 California
1665 Garden Avenue-U of M responsible? Shared?

Trees Pruned

Public

Grove Park-major deadwood in cottonwood over play structure
 1495 W. California-honey locust
 1390 Idaho-sugar maple storm damage/dieback
 1386 Idaho-storm damage in elm
 1434 Iowa -storm damage in green ash
 1427 California-storm damage in green ash
 1380 California-dead section in linden
 1376 California-storm damage in elm
 1489 Hoyt-storm damage in linden
 1881 Holton- honey locust
 1596 Vincent-hackberry, improved sightlines from driveway

Hazardous Trees on Private Property-Letters Sent out requesting action

1712 Tatum-storm damaged elm, suggest cleanup action taken
 1864 Asbury-split ash tree, suggest removal action taken
 1688 Arona-green ash storm damage, suggest pruning
 1852 Holton-storm damaged limb in green ash, suggest pruning

Trees Planted

1942 Summer Sugar Maple
 1984 Autumn Nortwoods Red Maple
 1790 Pascal Valley Forge Elm
 1547 Idaho Red Maple
 1433 Idaho Japanese Tree Lilac
 1423 Idaho Autumn Spire Red Maple
 2123 Hoyt Valley Forge Elm, tree planted last year but in severe decline
 planting contractor removed old tree
 1427 California Bur oak
 1427 Iowa Sugar Maple